


Nucleophilic 1,1-Difluoroethylation with Fluorinated Phosphonium Salt

Zuyong Deng,[†] Can Liu,^{†,‡} Xian-Liang Zeng,[†] Jin-Hong Lin,^{*,†,‡} and Ji-Chang Xiao^{*,†}[†]Key Laboratory of Organofluorine Chemistry, Shanghai Institute Of Organic Chemistry, University of Chinese Academy of Sciences, Chinese Academy of Sciences, 345 Lingling Road, Shanghai 200032, China[‡]School of Chemistry and Chemical Engineering, University of South China, Hengyang 421001, P. R. China

Supporting Information

ABSTRACT: The fluorinated phosphonium salt ($\text{Ph}_3\text{P}^+\text{CF}_2\text{CH}_3 \text{BF}_4^-$) was shown to act as a nucleophilic 1,1-difluoroethylation agent to enable difluoroethylation of aldehydes and imines.


The 1,1-difluoroethyl group (CF_2CH_3) has proved to be a valuable pharmacophore, as it not only is an isostere of a methoxy group but may also improve the bioactivity of the target molecules.¹ Consequently, significant efforts have been directed toward the exploration of general methods for the incorporation of this functionality. Traditional methods such as deoxyfluorination of carbonyl compounds suffer from harsh reaction conditions and a narrow substrate scope.² C–H fluorination is an operationally simple approach for the installation of the CF_2CH_3 moiety,³ and fluorination of terminal alkynes can also smoothly construct the CF_2CH_3 group.⁴ Direct difluoroethylation may serve as an efficient alternative to the fluorination strategies, but the studies in this area remains largely unexplored. Baran and co-workers disclosed that sodium difluoroethylsulfonate ($\text{CH}_3\text{CF}_2\text{SO}_2\text{Na}$) could be successfully applied to radical difluoroethylation reactions.⁵ Although $\text{CH}_3\text{CF}_2\text{Br}$ was found to be able to undergo difluoroethylation with aryl Grignard agents catalyzed by cobalt complexes, the high volatility of $\text{CH}_3\text{CF}_2\text{Br}$ results in operational inconveniences.⁶ Recently, it was found that $\text{TMSCF}_2\text{CH}_3$ was an effective nucleophilic difluoroethylation agent,⁷ but this agent is volatile, and its synthesis requires a multistep procedure. Therefore, the development of mild protocols for 1,1-difluoroethylation is still highly desirable.

Phosphonium salts have played an increasingly important role in a variety of research areas. Positively charged phosphorus increases the acidity of the adjacent C–H bond due to an inductive effect and can readily interact with a counteranion because of Coulombic interactions. Therefore, phosphonium salts have been widely used as ylide precursors,⁸ phase-transfer catalysts,⁹ Lewis acid catalysts,¹⁰ and so on. Apparently, these successful applications of phosphonium salts arise from the high electrophilicity of the cation. From this high electrophilicity it may be inferred that the cation or a substituent on the cation cannot act as a nucleophile to construct C–C bond. However, we recently discovered that a substituent on the phosphorus can be turned into a nucleophile, a process which was developed as a synthetic tool to carry out arylation¹¹ and difluoromethylation¹² reactions. The unprecedented findings opened up new

perspectives for the chemistry of phosphonium salts and prompted us to develop their applications further. We found that 1,1-difluoroethyl phosphonium salt ($\text{Ph}_3\text{P}^+\text{CF}_2\text{CH}_3 \text{BF}_4^-$) can be used as a nucleophilic difluoroethylation agent to enable difluoroethylation of aldehydes and imines.¹³ The preliminary results are described herein.

Phosphonium salt **2** ($\text{Ph}_3\text{P}^+\text{CF}_2\text{CH}_3 \text{BF}_4^-$) could be readily prepared from ethyl phosphonium salt **1** by a stepwise fluorination process (Scheme 1). Salt **2** is shelf-stable and could be purified simply by washing with organic solvents, allowing for its easy access.

Scheme 1. Preparation of 1,1-Difluoroethyl Phosphonium Salt 2




With a reliable approach for the synthesis of salt **2** established, optimization of difluoroethylation of aldehyde **3a** with salt **2** was conducted as shown in Table 1. As it has been previously shown that Cs_2CO_3 is a suitable promoter for the nucleophilic reactions with phosphonium salts,^{11,12} the effect of Cs_2CO_3 was first examined in our initial attempts at this difluoroethylation conversion. To our delight, the product was obtained by using THF as the solvent albeit in a low yield (Table 1, entry 1). A brief survey of the reaction solvents (Table 1, entries 2–6) revealed that a higher yield could be obtained in DMF (entry 6). Interestingly, the slow addition of a solution of salt **2** instead of adding the agent in one portion significantly increased the yield to 90% (entry 7 vs entry 6), which was similar to our previous observations.^{11,12} A series of other promoters were also screened, but no better results were obtained (Table 1, entries 8–11 vs 7). Increasing the reaction

Received: November 11, 2016

Published: December 3, 2016



Table 1. Screening Reaction Conditions^a


entry	solvent	promoter	4a, yield (%) ^b
1	THF	Cs ₂ CO ₃	28
2	cyclohexane	Cs ₂ CO ₃	4
3	toluene	Cs ₂ CO ₃	0
4	EtOAc	Cs ₂ CO ₃	23
5	CH ₃ CN	Cs ₂ CO ₃	8
6	DMF	Cs ₂ CO ₃	56
7 ^c	DMF	Cs ₂ CO ₃	90
8 ^c	DMF	K ₂ CO ₃	49
9 ^c	DMF	AcOCs	31
10 ^c	DMF	PhCO ₂ K	17
11 ^{c,d}	DMF	CsF	43
12 ^{c,e}	DMF	Cs ₂ CO ₃	90

^aConditions: **3a** (0.2 mmol), salt **2** (0.4 mmol), and promoter (2.5 equiv) in solvent at rt for 2 h. ^bDetermined by ¹⁹F NMR by using PhCF₃ as an internal standard. ^cA solution of salt **2** in DMF (1 mL) was added slowly into the solution of **3a** and promoter in DMF (1 mL) over 30 min. ^d5 equiv of CsF was used. ^eThe reaction scale was increased to 0.5 mmol (substrate **3a**).


scale did not lead to a decrease in the yield (Table 1, entry 12 vs 7).

With the optimized reaction conditions in hand (Table 1, entry 7), we then investigated the substrate scope for the nucleophilic difluoroethylation of carbonyls with salt **2**. As shown in Scheme 2, the examination of electronic effects showed that neither the electron-donating groups nor the electron-withdrawing groups had side effects for the conversion of aryl aldehydes (**4a–m**). The transformation was moderately


sensitive to steric effects, as evidenced by the moderate yield of **4e**. Heteroaryl aldehydes were also well tolerated and could be converted smoothly into the expected products in good yields (**4n–p**). Enolizable aldehyde was found to be inert toward difluoroethylation under these reaction conditions (**4q**). For the conversion of ketone, the yield was decreased dramatically (**4r**). Due to the lower reactivity of aliphatic aldehyde and ketone, the corresponding difluoroethylation products (**4q–r**) were obtained in very low yields. Attempts to increase their yields by reoptimizing the reaction conditions such as increasing the loading of reagent or elevating the reaction temperature were not successful.

The successful conversion of aryl aldehydes encouraged us to examine the difluoroethylation of imines. Compared with aldehydes, *N*-Ts imines exhibited lower reactivity and only moderate yields were obtained (Scheme 3). The substrates containing electron-donating groups could be well converted (**6a–f**). Although it may be expected that the aryl substrates containing electron-withdrawing group might be well transformed due to their higher electrophilicity, the difluoroethylation of 4-cyanophenylimine gave the desired product only in a very low yield (**6g**). That should be partially because the substrate would readily undergo hydrolysis with cesium carbonate under these reaction conditions.

On the basis of the above results and our previous studies toward nucleophilic reactions with phosphonium salts,^{11,12} the difluoroethylation conversion is proposed to occur through initial attack of Cs₂CO₃ at phosphonium salt **2** (Scheme 4). Since the attack of a nucleophile at the positively charged phosphorus would usually be along the axial direction and the subsequent pseudorotation would place the electronegative substituent in the other axial position,¹⁴ the trigonal bipyramidal phosphorus species **Int** would be readily generated from this nucleophilic addition process. Intermediate **Int** is highly reactive, and its decarboxylation would occur rapidly,


Scheme 2. Substrate Scope for the Difluoroethylation of Carbonyls^a

^aIsolated yields. ^bThe yields were determined by ¹⁹F NMR.

Scheme 3. Substrate Scope for the Difluoroethylation of Imines^a

^aIsolated yields. ^bThe yield was determined by ¹⁹F NMR.

Scheme 4. Proposed Reaction Mechanism



giving Ph_3PO and resulting in the cleavage of the $\text{P}-\text{CF}_2$ bond. The bond breaking may proceed concurrently with the nucleophilic addition of CH_3CF_2 group to the substrate. Hydrolysis gives the final product. Considering that the cyano group (in **4m** in Scheme 2) was compatible with the reaction conditions, we believe that the free CH_3CF_2^- ion may not be produced. The proposed mechanism was further supported by the almost full conversion of salt **2** into Ph_3PO as detected by ³¹P NMR spectroscopy and the observation of $\text{CH}_3\text{CF}_2\text{H}$ and $\text{CH}_3\text{CF}_2\text{CO}_2^-$ as determined by ¹⁹F NMR spectroscopy. $\text{CH}_3\text{CF}_2\text{H}$ and $\text{CH}_3\text{CF}_2\text{CO}_2^-$ should be formed via the nucleophilic attack of CH_3CF_2 moiety in **Int** at proton or CO_2 , respectively.

CONCLUSIONS

In summary, we have disclosed that 1,1-difluoroethyl phosphonium salt ($\text{Ph}_3\text{P}^+\text{CF}_2\text{CH}_3 \text{BF}_4^-$) can be successfully applied to 1,1-difluoroethylation of aldehydes and imines under mild conditions. This phosphonium salt is shelf stable and easy to access and, therefore, is reasonably expected to become an efficient 1,1-difluoroethylation agent. This work offers new opportunities to explore valuable applications of phosphonium salts in the chemistry of C–C bond construction. The strategy for nucleophilic fluoroalkylation with phosphonium salts may find synthetic utility in other research areas.

EXPERIMENTAL SECTION

All solvents were obtained from commercial sources and were extra dry grade. All glassware used was dried in a 120 °C oven and cooled in a desiccator before use. High-resolution mass data were recorded on a high-resolution mass spectrometer in EI or ESI mode. ¹H, ¹⁹F, ³¹P, and ¹³C NMR spectra were recorded on a 400 MHz or 500 MHz NMR spectrometer. ¹H NMR and ¹³C NMR chemical shifts were determined relative to internal $(\text{CH}_3)_4\text{Si}$ (TMS) at δ 0.0 ppm. ¹⁹F NMR chemical shifts were determined relative to internal CFCl_3 at 0.0 ppm. ³¹P NMR chemical shifts were determined relative to internal H_3PO_4 at 0.0 ppm.

Procedure for the Synthesis of 1,1-Difluoroethyl Phosphonium Salt. *Synthesis of Ethyl Phosphonium Salt 1.* Under N_2 atmosphere, the mixture of Ph_3P (8.5 g, 33 mmol), EtBr (3.3 g, 30 mmol), and *p*-xylene (20 mL) was stirred at 110 °C for 3 h. The mixture was cooled to room temperature. After filtration, the residue was washed with dry Et₂O (60 mL) and then dried under reduced pressure to afford the pure product.

*Ethyl triphenylphosphonium bromide (1):*¹⁵ 7.1 g, 63% yield, white solid; ¹H NMR (400 MHz, CDCl_3) δ 7.82–7.72 (m, 9H), 7.65 (td, J = 8.1, 3.5 Hz, 6H), 3.84–3.74 (m, 2H), 1.34 (dt, J = 20.1, 7.4 Hz, 3H); ³¹P NMR (162 MHz, CDCl_3) δ 26.28 (s, 1P).

Synthesis of 1-Fluoroethyl Phosphonium Salt 1'. Under argon atmosphere, into the mixture of **1** (3.9 g, 10 mmol) and THF (16 mL) at 0 °C was added a solution of BuLi (2.5 M, 11 mmol) in hexane (4.4 mL) dropwise via a syringe for 30 min. Upon completion of the addition, the resulting mixture was warmed to room temperature and was further stirred for 30 min to give the phosphonium ylide ($\text{Ph}_3\text{P}^+\text{CH}^-\text{Me}$). The solution of this ylide was stored in an ice bath before use.

Under argon atmosphere, into the solution of NFSI (*N*-fluoro-dibenzene-sulfonimide) (10.6 g, 33 mmol) in THF (30 mL) at 0 °C was added slowly the solution of the phosphonium ylide in 30 min. Upon completion of the addition, the resulting mixture was warmed to room temperature and was further stirred for 30 min. The reaction mixture was then poured into the aqueous solution of HBF_4 (1 M, 100 mL) and extracted with CH_2Cl_2 (50 mL \times 6). The combined organic phase was concentrated to afford a viscous liquid. Enough Et₂O (500 mL) was added into the liquid to precipitate the crude product, which was contaminated by NFSI, salt **1**, and salt **2**. After filtration, the solid was dissolved in CH_2Cl_2 (20 mL) to give a saturated solution. Enough Et₂O (500 mL) was added into the solution to precipitate the product. The solid was collected by filtration. The dissolving-precipitate process by $\text{CH}_2\text{Cl}_2/\text{Et}_2\text{O}$ was repeated once more to afford a solid. The solid was dissolved in CH_2Cl_2 (about 20 mL) to give a saturated solution. The solutions was stirred at 80 °C, and ethyl acetate (EA, 200 mL) was added slowly to precipitate the crude product. The crude product was collected by filtration. This dissolving-precipitate process by $\text{CH}_2\text{Cl}_2/\text{EA}$ was repeated for three to four times until the product was pure as detected by ³¹P NMR. The product was then dissolved in CH_2Cl_2 (80 mL), and the solution was then dried over anhydrous MgSO_4 . After filtration, the solvent was removed by concentration to give the pure product as a white solid (64%).

1-Fluoroethyl triphenylphosphonium tetrafluoroborate (1'): 2.67 g, 64% yield, white solid; mp 155–156 °C; ¹H NMR (400 MHz, CDCl_3) δ 7.83 (td, J = 7.2, 1.6 Hz, 3H), 7.78–7.67 (m, 12H), 6.98–6.78 (m, 1H), 1.91–1.72 (m, 3H); ¹⁹F NMR (376 MHz, CDCl_3) δ –152.89 (s, 1F), –152.95 (s, 3F), –193.71 (ddq, J = 69.6, 43.7, 25.9 Hz, 1F); ³¹P NMR (162 MHz, CDCl_3) δ 22.28 (d, J = 69.6 Hz, 1P); ¹³C{¹H} NMR (101 MHz, CDCl_3) δ 136.0 (d, J = 2.4 Hz), 134.2 (d, J = 9.7 Hz), 130.8 (d, J = 12.7 Hz), 114.7 (d, J = 84.2 Hz), 86.1 (dd, J =

191.7, 64.5 Hz), 16.4 (d, $J = 21.1$ Hz); IR (KBr) 3066, 2967, 1587, 1486, 1382, 1320, 1283, 1193, 1168, 1113, 1060, 997, 887, 754, 728, 711, 690, 541, 526, 498 cm^{-1} ; HRMS (ESI) calcd for $[\text{C}_{20}\text{H}_{19}\text{FP}]^+ [\text{M} - \text{BF}_4^-]^+$ 309.1203, found 309.1203.

Synthesis of 1,1-Difluoroethyl Phosphonium Salt 2. Under argon atmosphere, into the mixture of salt **1'** (3.96 g, 10 mmol) and THF (40 mL) at -78°C was added dropwise the solution of KHMDS [potassium bis(trimethylsilyl)amide] (1.0 M, 15 mmol) in THF (15 mL) via a syringe for 30 min. Upon completion of the addition, the resulting mixture was further stirred at 0°C until the solid disappeared (about 20 min) to give the fluorinated phosphonium ylide ($\text{Ph}_3\text{P}^+\text{CF}^-\text{Me}$). The solution of this ylide was unstable and needed to be stored at -78°C before use.

Under argon atmosphere, into the solution of NFSI (14.2 g, 45 mmol) in THF (40 mL) at 0°C was added the solution of the fluorinated ylide slowly in 30 min. Upon the completion of addition, the resulting mixture was warmed to room temperature and was further stirred for 10 min. The reaction mixture was then poured into the aqueous solution of HBF_4 (200 mL) and extracted with CH_2Cl_2 (50 mL \times 6). The combined organic phase was concentrated to afford a viscous liquid. Enough Et_2O (500 mL) was added into the liquid to precipitate the crude product, which was contaminated by NFSI and salt **1'**. After filtration under air atmosphere, the solid was dissolved in CH_2Cl_2 (about 20 mL) to give a saturated solution. Enough Et_2O (500 mL) was added into the solution to precipitate the product. The solid was collected by filtration under air atmosphere. The dissolving-precipitate process by $\text{CH}_2\text{Cl}_2/\text{Et}_2\text{O}$ was repeated once more to afford a solid. The solid was dissolved in CH_2Cl_2 (about 20 mL) to give a saturated solution. The solutions was stirred at 80°C , and EA (200 mL) was added slowly to precipitate the crude product. The crude product was collected by filtration. This dissolving-precipitate process by $\text{CH}_2\text{Cl}_2/\text{EA}$ was repeated three to four times until the product was pure as detected by ^{31}P NMR. The product was then dissolved in CH_2Cl_2 (100 mL), and the solution was then dried over anhydrous MgSO_4 . After filtration, the solvent was removed by concentration to give the pure product as a white solid (3.0 g, 72%).

1,1-Difluoroethyl triphenylphosphonium tetrafluoroborate (2): 3.0 g, 72% yield, white solid; mp $153\text{--}154^\circ\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ 7.93 (t, $J = 7.3$ Hz, 3H), 7.84–7.70 (m, 12H), 2.19 [td, $J = 21.8$ ($\text{CH}_3\text{--CF}_2$), 9.7 Hz, 3H]; ^{19}F NMR (376 MHz, CDCl_3) δ –86.85 (dq, $J = 95.8$, 21.8 Hz, 2F), –153.27 (s, 1F), –153.33 (s, 3F); ^{31}P NMR (162 MHz, CDCl_3) δ 25.15 (t, $J = 95.8$ Hz, 1P); $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, CDCl_3) δ 136.9 (d, $J = 3.2$ Hz), 134.6 (d, $J = 10.3$ Hz), 131.3 (d, $J = 13.1$ Hz), 124.0 (td, $J = 270.4$, 92.4 Hz), 112.3 (d, $J = 83.5$ Hz), 22.7 (dt, $J = 36.2$, 18.1 Hz); IR (KBr) 3069, 1587, 1485, 1441, 1389, 1284, 1142, 1109, 1055, 997, 936, 899, 753, 729, 689, 544, 526, 494 cm^{-1} ; HRMS (ESI) calcd for $[\text{C}_{20}\text{H}_{18}\text{F}_2\text{P}]^+ [\text{M} - \text{BF}_4^-]^+$ 327.1105, found 327.1105.

General Procedure for 1,1-Difluoroethylation of Aldehydes and Imines. In a glovebox, into a 25 mL Schlenk tube were added substrate (0.5 mmol), Cs_2CO_3 (407.3 mg, 1.25 mmol), and DMF (1 mL). The tube was sealed and then taken out from the glovebox. Into this mixture under argon atmosphere at room temperature, the solution of **2** (414 mg, 1.0 mmol) in DMF (2.0 mL) was added dropwise via a syringe for 30 min. Upon completion of the addition, the resulting mixture was further stirred at the same temperature for 10 min. The reaction was quenched by 3 N HCl (0.5 mL). The resulting mixture was diluted with water (40 mL) and extracted with DCM (4 \times 40 mL). The combined organic phase was dried over Na_2SO_4 . After filtration, the solvent was removed by concentration, and the residue was subjected to column chromatography (ethyl acetate and petroleum ether as the eluent) to afford the pure product.

1-([1,1'-Biphenyl]-4-yl)-2,2-difluoropropan-1-ol (4a): 106.3 mg, 86% yield, white solid; mp $89\text{--}90^\circ\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ 7.63–7.58 (m, 4H), 7.52 (d, $J = 8.0$ Hz, 2H), 7.45 (t, $J = 7.4$ Hz, 2H), 7.36 (t, $J = 7.4$ Hz, 1H), 4.91 (td, $J = 9.5$, 3.8 Hz, 1H), 2.48 (d, $J = 3.8$ Hz, 1H), 1.58 (t, $J = 18.8$ Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –100.66 ((ABq)qd, $\delta_{\text{AB}} = 267.0$ Hz, $J_{\text{AB}} = 241.9$ Hz, $J = 18.8$, 9.5 Hz, 2F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 141.6 (s), 140.6 (s), 135.6 (s), 128.9 (s), 127.7 (s), 127.5 (s), 127.2 (s, Hz), 127.1 (s), 123.4 (t, J

= 243.1 Hz), 75.6 (t, $J = 28.6$ Hz), 18.9 (t, $J = 26.3$ Hz); IR (KBr) 3612, 3457, 2922, 1489, 1449, 1394, 1232, 1186, 1126, 1058, 964, 924, 845, 802, 759, 738, 692, 610, 592, 524, 497 cm^{-1} ; HRMS (EI) calcd for $[\text{C}_{15}\text{H}_{14}\text{OF}_2]^+ [\text{M}^+]$ 248.1007, found 248.1008.

1-([1,1'-Biphenyl]-2-yl)-2,2-difluoropropan-1-ol (4b): 93.4 mg, 78% yield, colorless oil; mp $49\text{--}50^\circ\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ 7.72 (d, $J = 7.4$ Hz, 1H), 7.45–7.33 (m, 5H), 7.30 (d, $J = 7.3$ Hz, 2H), 7.25 (d, $J = 7.4$ Hz, 1H), 4.96 (dd, $J = 11.8$, 10.2 Hz, 1H), 2.13 (br, 1H), 1.42 (t, $J = 18.9$ Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –101.67 ((ABq)qd, $\delta_{\text{AB}} = 609.2$ Hz, $J_{\text{AB}} = 244.3$ Hz, $J = 18.8$, 10.2 Hz, 2F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 142.8 (s), 140.8 (s), 134.5 (s), 130.3 (s), 129.4 (s), 128.5 (s), 128.3 (s), 127.9 (s), 127.5 (s), 127.4 (s), 123.3 (t, $J = 244.0$ Hz), 71.8 (t, $J = 27.2$ Hz), 20.4 (t, $J = 26.4$ Hz); IR (KBr) 3430, 3060, 3026, 2930, 1597, 1480, 1439, 1389, 1285, 1236, 1184, 1129, 1110, 1057, 1047, 1009, 925, 878, 850, 775, 755, 704, 658, 612, 592, 573, 537 cm^{-1} ; HRMS (EI) calcd for $[\text{C}_{15}\text{H}_{14}\text{OF}_2]^+ [\text{M}^+]$ 248.1007, found 248.1012.

2,2-Difluoro-1-(4-methoxyphenyl)propan-1-ol (4c): 82.4 mg, 75% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ 7.35 (d, $J = 8.5$ Hz, 2H), 6.89 (d, $J = 8.5$ Hz, 2H), 4.78 (t, $J = 9.6$ Hz, 1H), 3.80 (s, 3H), 2.44 (br, 1H), 1.49 (t, $J = 18.9$ Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –101.11 (qd, $J = 18.9$, 9.6 Hz, 2F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 159.8 (s), 128.9 (s), 128.5 (s), 123.4 (t, $J = 242.9$ Hz), 113.8 (s), 75.4 (t, $J = 28.8$ Hz), 55.3 (s), 18.9 (t, $J = 26.4$ Hz); IR (KBr) 3462, 3005, 2920, 2841, 1614, 1587, 1516, 1465, 1444, 1391, 1305, 1251, 1177, 1127, 1068, 1032, 963, 924, 859, 839, 802, 789, 774, 645, 628, 616, 571, 520 cm^{-1} ; HRMS (EI) calcd for $[\text{C}_{10}\text{H}_{12}\text{O}_2\text{F}_2]^+ [\text{M}^+]$ 202.0800, found 202.0807.

2,2-Difluoro-1-(2-methoxyphenyl)propan-1-ol (4d): 71.0 mg, 60% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ 7.40 (d, $J = 7.6$ Hz, 1H), 7.31 (t, $J = 7.6$ Hz, 1H), 6.98 (t, $J = 7.6$ Hz, 1H), 6.91 (d, $J = 7.6$ Hz, 1H), 5.13 (td, $J = 11.1$, 6.7 Hz, 1H), 3.85 (s, 3H), 3.26 (d, $J = 6.7$ Hz, 1H), 1.57 (t, $J = 18.9$ Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –102.51 ((ABq)qd, $\delta_{\text{AB}} = 364.5$ Hz, $J_{\text{AB}} = 241.8$ Hz, $J = 18.9$, 11.1 Hz, 2F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 157.3 (s), 129.7 (s), 129.2 (s), 124.9 (s), 123.4 (t, $J = 244.1$ Hz), 120.9 (s), 111.1 (s), 75.4 (t, $J = 28.8$ Hz), 55.6 (s), 20.0 (t, $J = 26.4$ Hz); IR (KBr) 3444, 3006, 2945, 2841, 1603, 1589, 1494, 1465, 1441, 1390, 1289, 1245, 1185, 1128, 1050, 1026, 962, 924, 853, 814, 785, 757, 734, 658, 608, 581, 566, 506 cm^{-1} ; HRMS (EI) calcd for $[\text{C}_{10}\text{H}_{12}\text{O}_2\text{F}_2]^+ [\text{M}^+]$ 202.0800, found 202.0800.

2,2-Difluoro-1-mesitylpropan-1-ol (4e): 68.8 mg, 63% yield, white solid; mp $49\text{--}50^\circ\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ 6.86 (s, 2H), 5.31 (td, $J = 11.3$, 4.6 Hz, 1H), 2.43 (s, 6H), 2.28 (d, $J = 4.6$ Hz, 1H), 2.25 (s, 3H), 1.68 (t, $J = 18.9$ Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –98.31 ((ABq)qd, $\delta_{\text{AB}} = 614.4$ Hz, $J_{\text{AB}} = 245.4$ Hz, $J = 18.9$, 11.3 Hz, 2F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 137.8 (s), 129.6 (s), 124.40 (s), 124.39 (t, $J = 243.4$ Hz), 73.7 (dd, $J = 30.0$, 28.1 Hz), 21.3 (t, $J = 3.3$ Hz), 21.0 (s), 20.7 (s); IR (KBr) 3462, 3004, 2925, 1611, 1576, 1450, 1387, 1232, 1186, 1124, 1068, 1022, 954, 938, 917, 853, 821, 791, 727, 645, 635, 587, 522 cm^{-1} ; HRMS (EI) calcd for $[\text{C}_{12}\text{H}_{16}\text{OF}_2]^+ [\text{M}^+]$ 214.1164, found 214.1163.

2,2-Difluoro-1-(naphthalen-2-yl)propan-1-ol (4f): 86.6 mg, 78% yield, white solid; mp $53\text{--}54^\circ\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ 7.91 (s, 1H), 7.88–7.81 (m, 3H), 7.54 (d, $J = 8.6$ Hz, 1H), 7.52–7.46 (m, 2H), 5.02 (td, $J = 9.5$, 3.7 Hz, 1H), 2.55 (d, $J = 3.7$ Hz, 1H), 1.52 (t, $J = 18.9$ Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –100.18 to –100.41 (m, 2F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 134.2 (s), 133.5 (s), 133.0 (s), 128.2 (s), 128.1 (s), 127.7 (s), 126.7 (s), 126.5 (s), 126.4 (s), 124.8 (s), 123.5 (t, $J = 243.3$ Hz), 75.9 (t, $J = 28.8$ Hz), 19.0 (t, $J = 26.3$ Hz); IR (KBr) 3442, 3059, 3007, 2923, 1602, 1509, 1442, 1391, 1365, 1271, 1236, 1190, 1123, 1067, 963, 927, 886, 861, 824, 805, 758, 744, 654, 564, 545, 501, 480 cm^{-1} ; HRMS (EI) calcd for $[\text{C}_{13}\text{H}_{12}\text{OF}_2]^+ [\text{M}^+]$ 222.0851, found 222.0861.

2,2-Difluoro-1-(4-fluorophenyl)propan-1-ol (4g): 79.6 mg, 82% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ 7.41 (dd, $J = 8.3$, 5.6 Hz, 2H), 7.05 (t, $J = 8.3$ Hz, 2H), 4.82 (t, $J = 9.6$ Hz, 1H), 2.14 (s, 1H), 1.48 (t, $J = 18.9$ Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –100.93 ((ABq)qd, $\delta_{\text{AB}} = 219.1$ Hz, $J_{\text{AB}} = 247.8$ Hz, $J = 18.9$, 9.6 Hz, 2F), –113.35 to –113.47 (m, 1F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3)

δ 162.9 (d, J = 247.1 Hz), 132.4 (s), 129.0 (d, J = 8.2 Hz), 123.2 (t, J = 243.0 Hz), 115.3 (d, J = 21.6 Hz), 75.1 (t, J = 29.2 Hz), 18.7 (t, J = 26.3 Hz); IR (KBr) 3442, 2992, 1607, 1514, 1393, 1228, 1129, 1068, 927, 844, 804, 562 cm^{-1} ; HRMS (EI) calcd for $[\text{C}_9\text{H}_9\text{OF}_3] [\text{M}^+]$ 190.0600, found 190.0607.

1-(3-Chlorophenyl)-2,2-difluoropropan-1-ol (4h): 83.8 mg, 81% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ 7.45 (s, 1H), 7.35–7.25 (m, 3H), 4.81 (t, J = 9.2 Hz, 1H), 2.37 (br, 1H), 1.49 (t, J = 18.9 Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –100.42 ((ABq)qd, δ_{AB} = 356.60 Hz, J_{AB} = 249.1 Hz, J = 18.9, 9.2 Hz, 2F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 138.6 (d, J = 4.3 Hz), 134.4 (s), 129.6 (s), 128.8 (s), 127.4 (s), 125.5 (s), 123.1 (t, J = 242.7 Hz), 75.0 (dd, J = 28.7, 29.8 Hz), 18.7 (t, J = 26.2 Hz); IR (KBr) 3589, 3437, 3071, 3007, 2923, 1600, 1576, 1475, 1432, 1392, 1288, 1234, 1188, 1129, 1102, 1063, 964, 929, 819, 803, 776, 728, 707, 692, 656, 595, 514, 480 cm^{-1} ; HRMS (EI) calcd for $[\text{C}_9\text{H}_9\text{ClOF}_2] [\text{M}^+]$ 206.0305, found 206.0309.

1-(4-Bromophenyl)-2,2-difluoropropan-1-ol (4i): 95.4 mg, 76% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ 7.50 (d, J = 8.4 Hz, 2H), 7.31 (d, J = 8.4 Hz, 2H), 4.81 (t, J = 9.4 Hz, 1H), 2.47 (s, 1H), 1.48 (t, J = 18.9 Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –100.65 ((ABq)qd, δ_{AB} = 333.9 Hz, J_{AB} = 246.8 Hz, J = 18.9, 9.4 Hz, 2F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 135.6 (dd, J = 1.3, 4.4 Hz), 131.5 (s), 128.9 (t, J = 1.4 Hz), 123.1 (t, J = 244.0 Hz), 122.8 (s), 75.1 (dd, J = 28.4, 29.9 Hz), 18.6 (t, J = 26.3 Hz); IR (KBr) 3589, 3435, 3006, 2924, 2853, 1595, 1488, 1392, 1234, 1187, 1128, 1074, 1012, 964, 927, 849, 836, 794, 779, 634, 622, 599, 564, 551, 502 cm^{-1} ; HRMS (EI) calcd for $[\text{C}_9\text{H}_9\text{BrOF}_2] [\text{M}^+]$ 249.9799, found 249.9804.

2,2-Difluoro-1-(4-(trifluoromethyl)phenyl)propan-1-ol (4j): 82.6 mg, 68% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ 7.62 (d, J = 8.3 Hz, 2H), 7.56 (d, J = 8.2 Hz, 2H), 4.90 (t, J = 9.2 Hz, 1H), 2.68 (br, 1H), 1.48 (t, J = 18.9 Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –62.75 (s, 3F), –100.32 ((ABq)qd, δ_{AB} = 522.3 Hz, J_{AB} = 250.3 Hz, J = 18.9, 9.2 Hz, 2F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 140.5 (s), 130.9 (q, J = 32.5 Hz), 127.6 (s), 125.2 (q, J = 3.8 Hz), 124.0 (q, J = 273.4 Hz), 123.0 (t, J = 243.4 Hz), 75.1 (t, J = 29.4 Hz), 18.6 (t, J = 26.2 Hz); IR (KBr) 3608, 3447, 2924, 2361, 2341, 1622, 1420, 1394, 1327, 1234, 1167, 1126, 1068, 1019, 956, 929, 851, 788, 804, 766, 734, 679, 623, 609, 501 cm^{-1} ; HRMS (EI) calcd for $[\text{C}_{10}\text{H}_9\text{OF}_5] [\text{M}^+]$ 240.0568, found 240.0570.

2,2-Difluoro-1-(3-(trifluoromethyl)phenyl)propan-1-ol (4k): 88.5 mg, 73% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ 7.72 (s, 1H), 7.63 (d, J = 8.0 Hz, 1H), 7.61 (d, J = 8.0 Hz, 1H), 7.49 (t, J = 8.0 Hz, 1H), 4.91 (t, J = 9.0 Hz, 1H), 2.40 (br, 1H), 1.49 (t, J = 18.9 Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –62.71 (s, 3F), –100.47 ((ABq)qd, δ_{AB} = 532.2 Hz, J_{AB} = 248.4 Hz, J = 18.9, 9.0 Hz, 2F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 137.6 (d, J = 4.6 Hz), 130.8 (q, J = 32.5 Hz), 130.7 (s), 128.8 (s), 125.5 (q, J = 3.7 Hz), 124.1 (q, J = 3.8 Hz), 124.0 (q, J = 272.4 Hz), 123.0 (t, J = 244.4 Hz), 75.1 (dd, J = 30.1, 29.0 Hz), 18.6 (t, J = 26.2 Hz); IR (KBr) 3603, 3439, 3011, 2923, 1619, 1494, 1451, 1394, 1330, 1234, 1167, 1127, 1075, 1003, 965, 928, 813, 193, 758, 704, 683, 654, 613, 613, 590, 555, 506 cm^{-1} ; HRMS (EI) calcd for $[\text{C}_{10}\text{H}_9\text{OF}_5] [\text{M}^+]$ 240.0568, found 240.0578.

2,2-Difluoro-1-(2-(trifluoromethyl)phenyl)propan-1-ol (4l): 98.1 mg, 81% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ 7.86 (d, J = 7.8 Hz, 1H), 7.67 (d, J = 7.8 Hz, 1H), 7.61 (t, J = 7.8 Hz, 1H), 7.46 (t, J = 7.8 Hz, 1H), 5.25 (t, J = 10.5 Hz, 1H), 2.18 (br, 1H), 1.57 (t, J = 18.9 Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –57.21 (t, J = 4.0 Hz, 3F), –101.00 to –102.05 (m, 1F), –102.59 to –103.73 (m, 1F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 135.6 (s), 132.2 (s), 129.5 (t, J = 2.9 Hz), 128.9 (q, J = 30.1 Hz), 128.9 (s), 125.8 (q, J = 5.8 Hz), 124.1 (q, J = 274.5 Hz), 122.5 (t, J = 244.5 Hz), 70.8 (tq, J = 27.5, 2.6 Hz), 20.6 (t, J = 26.4 Hz); IR (KBr) 3604, 3433, 3011, 1894, 1609, 1588, 1457, 1392, 1313, 1235, 1167, 1123, 1070, 1054, 1035, 964, 931, 855, 817, 771, 750, 677, 678, 613, 599 cm^{-1} ; HRMS (EI) calcd for $[\text{C}_{10}\text{H}_9\text{OF}_5] [\text{M}^+]$ 240.0568, found 240.0576.

3-(2,2-Difluoro-1-hydroxypropyl)benzonitrile (4m): 66.0 mg, 69% yield, yellow oil; ^1H NMR (400 MHz, CDCl_3) δ 7.75 (s, 1H), 7.68 (d, J = 7.8 Hz, 1H), 7.60 (d, J = 7.8 Hz, 1H), 7.46 (t, J = 7.8 Hz, 1H), 4.86 (t, J = 9.1 Hz, 1H), 3.14 (br, 1H), 1.48 (t, J = 18.9 Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –98.56 to –99.48 (m, 1F), –101.04 to –101.99

(m, 1F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 138.5 (s), 132.2 (s), 131.9 (s), 131.0 (s), 129.1 (s), 122.9 (t, J = 243.6 Hz), 118.6 (s), 112.3 (s), 74.5 (t, J = 29.9 Hz), 18.7 (t, J = 26.1 Hz); IR (KBr) 3453, 3007, 2234, 1585, 1484, 1438, 1391, 1275, 1259, 1235, 1190, 1141, 1066, 966, 927, 787, 750, 692, 613, 588, 538 cm^{-1} ; HRMS (EI) calcd for $[\text{C}_{10}\text{H}_9\text{ONF}_2] [\text{M}^+]$ 197.0647, found 197.0654.

2,2-Difluoro-1-(quinolin-3-yl)propan-1-ol (4n): 82.6 mg, 77% yield, white solid; mp 126–127 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ 8.92 (s, 1H), 8.27 (s, 1H), 8.10 (d, J = 8.5 Hz, 1H), 7.83 (d, J = 8.2 Hz, 1H), 7.72 (t, J = 7.7 Hz, 1H), 7.56 (t, J = 7.6 Hz, 1H), 5.08 (t, J = 9.2 Hz, 1H), 3.49 (br, 1H), 1.57 (t, J = 18.9 Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –100.30 ((ABq)qd, δ_{AB} = 747.1 Hz, J_{AB} = 251.2 Hz, J = 18.9, 9.2 Hz, 2F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 149.5 (s), 147.4 (s), 135.4 (s), 130.8 (s), 130.1 (s), 128.4 (s), 128.1 (s), 127.6 (s), 127.1 (s), 123.3 (t, J = 243.4 Hz), 73.5 (t, J = 30.3 Hz), 19.0 (t, J = 26.2 Hz); IR (KBr) 3061, 2958, 1991, 1652, 1595, 1391, 1260, 1238, 1187, 1123, 1091, 1014, 964, 919, 818, 775, 758, 616, 593, 550 cm^{-1} ; HRMS (EI) calcd for $[\text{C}_{12}\text{H}_{11}\text{NOF}_2] [\text{M}^+]$ 223.0803, found 223.0811.

1-(Benzofuran-2-yl)-2,2-difluoropropan-1-ol (4o): 90.9 mg, 82% yield, yellow solid; mp 65–66 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ 7.82 (d, J = 8.3 Hz, 1H), 7.75 (dd, J = 6.5, 2.5 Hz, 1H), 7.40–7.29 (m, 3H), 5.16 (t, J = 9.1 Hz, 1H), 2.60 (br, 1H), 1.63 (t, J = 18.9 Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –100.59 ((ABq)qd, δ_{AB} = 349.4 Hz, J_{AB} = 249.9 Hz, J = 18.9, 9.1 Hz, 2F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 140.0 (s), 139.8 (s), 139.2 (s), 124.7 (s), 124.5 (s), 123.8 (s), 123.2 (s), 122.8 (t, J = 244.3 Hz), 122.4 (s), 73.0 (t, J = 30.6 Hz), 19.0 (t, J = 25.9 Hz); IR (KBr) 3430, 3059, 3000.5, 2924, 1458, 1437, 1391, 1334, 1308, 1260, 1234, 1186, 1126, 1046, 927, 862, 839, 804, 747, 726, 632, 609, 562 cm^{-1} ; HRMS (EI) calcd for $[\text{C}_{11}\text{H}_{10}\text{OF}_2] [\text{M}^+]$ 228.0415, found 228.0421.

1-(Benzofuran-2-yl)-2,2-difluoropropan-1-ol (4p): 81.4 mg, 76% yield, colorless oil; ^1H NMR (400 MHz, CDCl_3) δ 7.56 (d, J = 7.6 Hz, 1H), 7.48 (d, J = 7.6 Hz, 1H), 7.30 (t, J = 7.6 Hz, 1H), 7.23 (t, J = 7.6 Hz, 1H), 6.82 (s, 1H), 4.97 (t, J = 9.4 Hz, 1H), 2.56 (br, 1H), 1.71 (t, J = 18.8 Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –99.89 to –100.84 (m, 1F), –100.95 to –101.88 (m, 1F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 154.9 (s), 152.3 (s), 127.7 (s), 124.9 (s), 123.2 (s), 122.2 (t, J = 244.4 Hz), 121.4 (s), 111.5 (s), 106.1 (s), 70.9 (t, J = 31.0 Hz), 19.7 (t, J = 26.0 Hz); IR (KBr) 3428, 2992, 1476, 1454, 1393, 1238, 1138, 1061, 1009, 931, 885, 805, 751 cm^{-1} ; HRMS (EI) calcd for $[\text{C}_{11}\text{H}_{10}\text{O}_2\text{F}_2] [\text{M}^+]$ 212.0643, found 212.0644.

N-(1-(1,1'-Biphenyl-4-yl)-2,2-difluoropropyl)-4-methylbenzenesulfonamide (6a): 130.6 mg, 66% yield, white solid; mp 188–189 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ 7.52–7.31 (m, 9H), 7.12 (d, J = 8.0 Hz, 2H), 7.05 (d, J = 8.0 Hz, 2H), 5.27 (d, J = 8.5 Hz, 1H), 4.59 (td, J = 13.3, 8.5 Hz, 1H), 2.27 (s, 3H), 1.61 (t, J = 18.7 Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –98.35 to –101.08 (m, 2F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 143.4 (s), 141.4 (s), 140.3 (s), 137.2 (s), 133.1 (s), 129.3 (s), 128.8 (s), 128.6 (s), 127.6 (s), 127.1 (s), 127.0 (s), 122.3 (t, J = 245.8 Hz), 62.1 (t, J = 26.8 Hz), 21.7 (t, J = 26.4 Hz), 21.4 (s); IR (KBr) 3278, 3035, 2972, 2925, 1920, 1600, 1568, 1489, 1451, 1410, 1389, 1306, 1236, 1192, 1151, 1118, 1083, 1021, 1006, 963, 949, 911, 864, 853, 812, 763, 739, 696, 670, 611.571, 552, 544, 511 cm^{-1} ; HRMS (ESI) calcd for $[\text{C}_{22}\text{H}_{22}\text{O}_2\text{NSF}_2] [\text{M} + \text{H}^+]$ 402.1333, found 402.1334.

N-(2,2-Difluoro-1-(p-tolyl)propyl)-4-methylbenzenesulfonamide (6b): 93.2 mg, 55% yield, white solid; mp 196–197 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ 7.49 (d, J = 8.2 Hz, 2H), 7.07 (d, J = 8.2 Hz, 2H), 6.97 (d, J = 8.4 Hz, 2H), 6.94 (d, J = 8.4 Hz, 2H), 5.24 (d, J = 8.0 Hz, 1H), 4.49 (td, J = 13.0, 8.0 Hz, 1H), 2.32 (s, 3H), 2.26 (s, 3H), 1.54 (t, J = 18.7 Hz, 3H); ^{19}F NMR (376 MHz, CDCl_3) δ –99.71 ((ABq)qd, δ_{AB} = 613.7 Hz, J_{AB} = 244.3 Hz, J = 18.7, 13.0 Hz, 2F); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 143.3 (s), 138.4 (s), 137.3 (s), 131.3 (s), 129.3 (s), 129.1 (s), 128.0 (s), 127.1 (s), 122.3 (t, J = 245.0 Hz), 62.1 (dd, J = 25.6, 27.4 Hz), 21.6 (t, J = 26.3 Hz), 21.4 (s), 21.1 (s); IR (KBr) 3269, 3041, 3008, 2981, 1922, 1741, 1598, 1519, 1496, 1455, 1389, 1331, 1308, 1287, 1236, 1213, 1194, 1165, 1128, 1090, 1025, 961, 937, 911, 846, 813, 802, 781, 721, 705, 670, 645, 627, 582, 559, 545 cm^{-1} ; HRMS (ESI) calcd for $[\text{C}_{17}\text{H}_{20}\text{O}_2\text{NSF}_2] [\text{M} + \text{H}^+]$ 340.1177, found 340.1177.

N-(2,2-Difluoro-1-(4-methoxyphenyl)propyl)-4-methylbenzenesulfonamide (**6c**): 96.0 mg, 54% yield, white solid; mp 156–157 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.49 (d, *J* = 8.2 Hz, 2H), 7.08 (d, *J* = 8.2 Hz, 2H), 6.99 (d, *J* = 8.6 Hz, 2H), 6.68 (d, *J* = 8.6 Hz, 2H), 5.32 (d, *J* = 8.8 Hz, 1H), 4.49 (td, *J* = 12.6, 8.8 Hz, 1H), 3.74 (s, 3H), 2.32 (s, 3H), 1.54 (t, *J* = 18.6 Hz, 3H); ¹⁹F NMR (376 MHz, CDCl₃) δ –99.84 ((ABq)qd, δ_{AB} = 618.3 Hz, J_{AB} = 246.6 Hz, *J* = 18.6, 12.6 Hz, 2F); ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 159.6 (s), 143.2 (s), 137.3 (s), 129.31 (s), 129.28 (s), 127.0 (s), 126.4 (s), 122.4 (t, *J* = 243.9 Hz), 113.8 (s), 61.8 (dd, *J* = 25.8, 27.9 Hz), 55.2 (s), 21.5 (t, *J* = 26.3 Hz), 21.4 (s); IR (KBr) 3279, 3066, 3010, 2978, 2939, 2844, 1616, 1599, 1585, 1519, 1496, 1457, 1390, 1323, 1320, 1285, 1259, 1233, 1198, 1185, 1164, 1128, 1090, 1035, 956, 940, 912, 868, 845, 812, 727, 705, 670, 642, 626, 580, 559, 547 cm^{–1}; HRMS (ESI) calcd for [C₁₇H₂₀O₃NSF₂] [M + H⁺] 356.1126, found 356.1126.

N-(2,2-Difluoro-1-mesitylpropyl)-4-methylbenzenesulfonamide (**6d**): 92.1 mg, 50% yield, white solid; mp 160–161 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.41 (d, *J* = 8.3 Hz, 2H), 7.00 (d, *J* = 8.3 Hz, 2H), 6.73 (s, 1H), 6.56 (s, 1H), 5.31 (d, *J* = 9.2 Hz, 1H), 5.24–5.06 (m, 1H), 2.30 (s, 3H), 2.29 (s, 3H), 2.17 (s, 6H), 1.63 (t, *J* = 18.7 Hz, 3H); ¹⁹F NMR (376 MHz, CDCl₃) δ –93.22 to –94.16 (m, 1F), –94.43 to –95.35 (m, 1F); ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 143.2 (s), 137.8 (s), 137.7 (s), 137.0 (s), 136.3 (s), 131.4 (s), 129.3 (s), 129.1 (s), 127.7 (d, *J* = 2.4 Hz), 126.6 (s), 123.4 (t, *J* = 245.1 Hz), 58.4 (dd, *J* = 29.5, 26.0 Hz), 22.2 (t, *J* = 26.9 Hz), 21.4 (s), 21.2 (s), 21.1 (t, *J* = 5.9 Hz), 20.7 (s); IR (KBr) 3335, 2971, 1613, 1600, 1577, 1498, 1441, 1386, 1336, 1289, 1230, 1189, 1168, 1148, 1084, 1028, 971, 948, 919, 903, 857, 835, 819, 805, 726, 707, 670, 629, 595, 573, 549 cm^{–1}; HRMS (ESI) calcd for [C₁₉H₂₄O₂NSF₂] [M + H⁺] 368.1490, found 368.1490.

N-(2,2-Difluoro-1-phenylpropyl)-4-methylbenzenesulfonamide (**6e**): 88.3 mg, 54% yield, white solid; mp 175–176 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.49 (d, *J* = 8.3 Hz, 2H), 7.24–7.14 (m, 3H), 7.08–7.04 (m, 4H), 5.24 (d, *J* = 8.9 Hz, 1H), 4.64–4.44 (m, 1H), 2.31 (s, 3H), 1.56 (t, *J* = 18.7 Hz, 3H); ¹⁹F NMR (376 MHz, CDCl₃) δ –99.84 ((ABq)qd, δ_{AB} = 522.9 Hz, J_{AB} = 244.8 Hz, *J* = 18.7, 13.5 Hz, 2F); ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 143.3 (s), 137.2 (s), 134.3 (d, *J* = 2.7 Hz), 129.3 (s), 128.5 (s), 128.4 (s), 128.1 (s), 127.0 (s), 122.3 (t, *J* = 245.0 Hz), 62.3 (dd, *J* = 27.7, 25.4 Hz), 21.6 (t, *J* = 26.4 Hz), 21.4 (s); IR (KBr) 3266, 3069, 3013, 2974, 1600, 1497, 1459, 1448, 1389, 1332, 1309, 1269, 1234, 1190, 1166, 1127, 1095, 1070, 936, 953, 911, 861, 831, 813, 752, 703, 676, 643, 580, 558, 548 cm^{–1}; HRMS (ESI) calcd For [C₁₆H₁₈O₂NSF₂] [M + H⁺] 326.1020, found 326.1021.

N-(2,2-Difluoro-1-(naphthalen-2-yl)propyl)-4-methylbenzenesulfonamide (**6f**): 74.2 mg, 39% yield, white solid; mp 134–135 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.78–7.72 (m, 1H), 7.67–7.61 (m, 2H), 7.49–7.38 (m, 5H), 7.20 (d, *J* = 8.4 Hz, 1H), 6.86 (d, *J* = 8.0 Hz, 2H), 5.41 (d, *J* = 6.5 Hz, 1H), 4.74–4.63 (m, 1H), 2.09 (s, 3H), 1.61 (t, *J* = 18.7 Hz, 3H); ¹⁹F NMR (376 MHz, CDCl₃) δ –99.42 ((ABq)qd, δ_{AB} = 516.3 Hz, J_{AB} = 244.6 Hz, *J* = 18.7, 13.0 Hz, 2F); ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 143.3 (s), 137.0 (s), 133.0 (s), 132.8 (s), 131.3 (s), 129.1 (s), 128.4 (s), 128.2 (s), 128.0 (s), 127.5 (s), 127.0 (s), 126.5 (s), 126.3 (s), 125.0 (s), 122.4 (t, *J* = 245.7 Hz), 62.6 (dd, *J* = 27.9, 25.7 Hz), 21.7 (t, *J* = 26.2 Hz), 21.2 (s); IR (KBr) 3273, 3062, 3007, 2953, 1921, 1594, 1511, 1492, 1452, 1432, 1395, 1335, 1308, 1270, 1247, 1202, 1165, 1126, 1086, 1040, 1017, 975, 936, 910, 866, 816, 806, 781, 766, 752, 705, 670, 603, 564, 543 cm^{–1}; HRMS (ESI) calcd for [C₂₀H₂₀O₂NSF₂] [M + H⁺] 376.1177, found 376.1175. Anal. Calcd for C₂₀H₁₉O₂NSF₂: C, 63.98; H, 5.10; N, 3.73; F, 10.12; S, 8.54. Found: C, 63.60; H, 5.13; N, 3.54; F, 10.34; S, 8.59.

N-(1-(4-Cyanophenyl)-2,2-difluoropropyl)-4-methylbenzenesulfonamide (**6g**): 16% yield determined by ¹⁹F NMR. In order to isolate this product to get the full characterization data, the reaction scale was increased (1.25 mmol of substrate): 54.6 mg, 12%, white solid; mp 152–153 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.53 (d, *J* = 8.1 Hz, 2H), 7.46 (d, *J* = 8.1 Hz, 2H), 7.25 (d, *J* = 8.3 Hz, 2H), 7.11 (d, *J* = 8.3 Hz, 2H), 6.11 (d, *J* = 9.2 Hz, 1H), 4.63 (dt, *J* = 15.1, 9.2 Hz, 1H), 2.36 (s, 3H), 1.61 (t, *J* = 18.3 Hz, 3H); ¹⁹F NMR (376 MHz, CDCl₃) δ –97.74 to –100.50 (m, 2F); ¹³C NMR (101 MHz, CDCl₃) δ 144.0

(s), 139.3 (d, *J* = 1.7 Hz), 136.7 (s), 132.0 (s), 129.5 (s), 129.1 (s), 126.9 (s), 121.8 (t, *J* = 249.7 Hz), 118.2 (s), 112.3 (s), 61.8 (dd, *J* = 29.6, 25.8 Hz), 21.47 (t, *J* = 26.2 Hz), 21.44 (s); IR (neat) ν = 3271, 1418, 1330, 1235, 1184, 1164, 1089, 911, 668, 585 cm^{–1}; HRMS (ESI) calcd for C₁₇H₁₅O₂N₂F₂S [M – H][–] 349.0822, found 349.0828.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.6b02723.

¹H, ¹⁹F, ³¹P, and ¹³C NMR spectra of products (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: jlin@sioc.ac.cn. Fax: +86-21-6416-6128. Tel: +86-21-5492-5541.

*E-mail: jchxiao@sioc.ac.cn. Fax: +86-21-6416-6128. Tel: +86-21-5492-5340.

ORCID

Jin-Hong Lin: 0000-0002-7000-9540

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank National Basic Research Program of China (2015CB931900, 2012CBA01200), the National Natural Science Foundation (21421002, 21472222, 21502214, 21672242), the Chinese Academy of Sciences (XDA02020105, XDA02020106), and the Science and Technology Commission of Shanghai Municipality (15DZ1200102, 14ZR1448800) for financial support.

■ REFERENCES

- (a) Coteron, J. M.; Marco, M.; Esquivias, J.; Deng, X.; White, K. L.; White, J.; Koltun, M.; El Mazouni, F.; Kokkonda, S.; Katneni, K.; Bhamidipati, R.; Shackelford, D. M.; Angulo-Barturen, I.; Ferrer, S. B.; Jiménez-Díaz, M. B.; Gamo, F.-J.; Goldsmith, E. J.; Charman, W. N.; Bathurst, I.; Floyd, D.; Matthews, D.; Burrows, J. N.; Rathod, P. K.; Charman, S. A.; Phillips, M. A. *J. Med. Chem.* **2011**, *54*, 5540–5561. (b) Anderson, M. O.; Zhang, J.; Liu, Y.; Yao, C.; Phuan, P.-W.; Verkman, A. S. *J. Med. Chem.* **2012**, *55*, 5942–5950. (c) Liu, Y.; Esteva-Font, C.; Yao, C.; Phuan, P. W.; Verkman, A. S.; Anderson, M. O. *Bioorg. Med. Chem. Lett.* **2013**, *23*, 3338–3341.
- (a) Markovskij, L. N.; Pashinnik, V. E.; Kirsanov, A. V. *Synthesis* **1973**, 1973, 787–789. (b) Middleton, W. J. *J. Org. Chem.* **1975**, *40*, 574–578. (c) York, C.; Surya Prakash, G. K.; Olah, G. A. *Tetrahedron* **1996**, *52*, 9–14.
- (a) Xia, J.-B.; Zhu, C.; Chen, C. *J. Am. Chem. Soc.* **2013**, *135*, 17494–17500. (b) Xu, P.; Guo, S.; Wang, L.; Tang, P. *Angew. Chem., Int. Ed.* **2014**, *53*, 5955–5958.
- Okoromoba, O. E.; Han, J.; Hammond, G. B.; Xu, B. *J. Am. Chem. Soc.* **2014**, *136*, 14381–14384.
- Zhou, Q.; Ruffoni, A.; Gianatassio, R.; Fujiwara, Y.; Sella, E.; Shabat, D.; Baran, P. S. *Angew. Chem., Int. Ed.* **2013**, *52*, 3949–3952.
- Ohtsuka, Y.; Yamakawa, T. *J. Fluorine Chem.* **2016**, *185*, 96–102.
- (a) Mogi, R.; Morisaki, K.; Hu, J.; Prakash, G. K. S.; Olah, G. A. *J. Fluorine Chem.* **2007**, *128*, 1098–1103. (b) Li, X.; Zhao, J.; Wang, Y.; Rong, J.; Hu, M.; Chen, D.; Xiao, P.; Ni, C.; Wang, L.; Hu, J. *Chem. - Asian J.* **2016**, *11*, 1789–1792.
- (a) Murphy, P. J.; Brennan, J. *Chem. Soc. Rev.* **1988**, *17*, 1–30. (b) Maryanoff, B. E.; Reitz, A. B. *Chem. Rev.* **1989**, *89*, 863–927. (c) Marsden, S. P. *Nat. Chem.* **2009**, *1*, 685–687. (d) Byrne, P. A.; Gilheany, D. G. *Chem. Soc. Rev.* **2013**, *42*, 6670–6696.
- For recent reviews, see: (a) Werner, T. *Adv. Synth. Catal.* **2009**, *351*, 1469–1481. (b) Enders, D.; Nguyen, T. V. *Org. Biomol. Chem.*

2012, 10, 5327–5331. (c) Liu, S.; Kumatabara, Y.; Shirakawa, S. *Green Chem.* **2016**, 18, 331–341. For recent examples, see: (d) He, R.; Ding, C.; Maruoka, K. *Angew. Chem., Int. Ed.* **2009**, 48, 4559–4561. (e) Shirakawa, S.; Kasai, A.; Tokuda, T.; Maruoka, K. *Chem. Sci.* **2013**, 4, 2248–2252. (f) Cao, D.; Zhang, J.; Wang, H.; Zhao, G. *Chem. - Eur. J.* **2015**, 21, 9998–10002. (g) Ge, L.; Lu, X.; Cheng, C.; Chen, J.; Cao, W.; Wu, X.; Zhao, G. *J. Org. Chem.* **2016**, 81, 9315–9325.

(10) For recent reviews, see: (a) Bayne, J. M.; Stephan, D. W. *Chem. Soc. Rev.* **2016**, 45, 765–774. For recent examples, see: (b) Caputo, C. B.; Hounjet, L. J.; Dobrovetsky, R.; Stephan, D. W. *Science* **2013**, 341, 1374–1377. (c) Perez, M.; Hounjet, L. J.; Caputo, C. B.; Dobrovetsky, R.; Stephan, D. W. *J. Am. Chem. Soc.* **2013**, 135, 18308–18310. (d) Mehta, M.; Holthausen, M. H.; Mallov, I.; Perez, M.; Stephan, D. W.; Qu, Z.-W.; Grimme, S. *Angew. Chem., Int. Ed.* **2015**, 54, 8250–8254. (e) Zhu, J.; Perez, M.; Caputo, C. B.; Stephan, D. W. *Angew. Chem., Int. Ed.* **2016**, 55, 1417–1421.

(11) Deng, Z.; Lin, J.-H.; Xiao, J.-C. *Nat. Commun.* **2016**, 7, 10337.

(12) Deng, Z.; Lin, J.-H.; Cai, J.; Xiao, J.-C. *Org. Lett.* **2016**, 18, 3206–3209.

(13) For recent studies on the use of fluorinated phosphonium salts as *gem*-difluoroalkylation agents, see: (a) Levin, V. V.; Trifonov, A. L.; Zemtsov, A. A.; Struchkova, M. I.; Arkhipov, D. E.; Dilman, A. D. *Org. Lett.* **2014**, 16, 6256–6259. (b) Panferova, L. I.; Tsymbal, A. V.; Levin, V. V.; Struchkova, M. I.; Dilman, A. D. *Org. Lett.* **2016**, 18, 996–999. (c) Lin, Q.-Y.; Ran, Y.; Xu, X.-H.; Qing, F.-L. *Org. Lett.* **2016**, 18, 2419–2422. (d) Ran, Y.; Lin, Q.-Y.; Xu, X.-H.; Qing, F.-L. *J. Org. Chem.* **2016**, 81, 7001–7007.

(14) (a) Cristau, H. J.; Plenat, F. *The Chemistry of Organophosphorus Compounds: Phosphonium Salts, Ylides and Phosphoranes*; Hartley, F. R., Ed.; John Wiley & Sons: Chichester, 1994; Vol. 3, pp 45–184. (b) Albright, T. A.; Burdett, J. K.; Whangbo, M.-H. *Orbital Interactions in Chemistry*; John Wiley & Sons: Chichester, 2013; pp 359–400. (c) Allen, D. W. *J. Chem. Soc. B* **1970**, 1490–1493. (d) Byrne, P. A.; Ortin, Y.; Gilheany, D. G. *Chem. Commun.* **2015**, 51, 1147–1150.

(15) Byrne, P. A.; Gilheany, D. G. *J. Am. Chem. Soc.* **2012**, 134, 9225–9239.